

## **Proprioceptive Neuromuscular Facilitation (PNF I) Bibliography**

Recommended:

Valenza, M.C., Torres-Sanchez, I., Cabrera-Martos, I, Valenza-Demet, G., Cano-Cappellacci, M, 2016 “Acute Effects of Contract-Relax Stretching vs. TENS in Young Subjects with Anterior Knee Pain: A Randomized Controlled Trial” *The Journal of Strength and Conditioning Research*, 30(8), 2271-8.

Yildirim, M.S., Ozyurek, S., Tosun, O., Uzer, S., Gelecek, N., 2016 “Comparison of effects of static, proprioceptive neuromuscular facilitation and Mulligan stretching on hip flexion range of motion: a randomized controlled trial” *National Center for Biotechnology Information*, 33(1): 89-94.

Hoogenboom, B.J., 2015 “Rolling Revisited: Using rolling to assess and treat neuromuscular control and coordination of the core and extremities of athletes” *International Journal of Sports Physical Therapy*, 10(6): 787-802.

Chow, E., 2015 “The longitudinal impact of parent distress and behavior on functional outcomes among youth with chronic pain” *The Journal of Pain*, Boston University Theses & Dissertations

Kent, P., Laird, R., Haines, T., 2015 “The effect of changing movement and posture using motion-sensor biofeedback, versus guidelines-based care, on the clinical outcomes of people with sub acute pr chronic low back pain- a multicenter, cluster-randomised, placebo-controlled pilot trial” *BMC Musculoskeletal Disorder*, 16(131): 1-19.

Stanton, T.R., Leake, H.B., Chalmers, K.J., Moseley, G.L., 2015 “Evidence of Impaired Proprioception in Chronic, Idiopathic Neck Pain: Systematic Review and Meta-Analysis” *Research Report*, 96(6): 877-887.

Lee, B.K., 2015 “Effects of the combined PNF and deep breathing exercises on the ROM and the VAS score of a frozen shoulder patient: Single case study” *Journal of Exercise Rehabilitation*, 11(5): 276-281.

Park, K., Seo, K., 2014 “The Effects on the Pain Index and Lumbar Flexibility of Obese Patients with Low Back Pain after PNF Scapular and PNF Pelvic Patterns” *Journal of Physical Therapy Science*, 26(10), 1571-4.

Mitsuo, A., 2014 “Effect of remote after-effects of resistive static contraction of the pelvic depressors on improvement of restricted wrist flexion range of motion in patients with restricted wrist flexion range of motion” *Journal of Bodywork & Movement Therapies*, 19(3), 442-6.

Ribeiro, T.S., Gomes, E.M., Sousa Silva, W.H., Alencar Caldas, V.V., Araujo Silva, D.L., Costa Cavalcanti, F.A., Lindquist, A.R. 2014 “Effects of a training program based on the Proprioceptive Neuromuscular Facilitation method on post-stroke moroe recovery: A preliminary study” *Journal of Bodywork and Movement Therapies*, 18: 526-532.

Voight, M.L., Hoogenboom, B.J., 2008 “The Chop and Lift Reconsidered: Integrating Neuromuscular Priciples into Orthopedic and Sports Rehabilitation” *N Am J Sports Phys Therapy*, 3(3):151-159.

PNF in Practice, Adler, Susan, Beckers, Dominiek, and Buck, Math. 2<sup>nd</sup> Ed., NY: Springer Verlag, 2000, Ch. 2, 3, 6, 11, & 12. Available on [www.amazon.com](http://www.amazon.com).

Rational Manual Therapies, Basmajian and Nyberg, Chapter 11 “Proprioceptive Neuromuscular Facilitation” by Vicky Saliba, Gregory S. Johnson and Cheryl Wardlaw. Available online at [www.instituteofphysicalart.com](http://www.instituteofphysicalart.com) or we will mail at cost \$5.

Andersen L.L., Magnusson S.P., Nielsen M., Haleem J., Poulsen K., Aagaard P. Neuromuscular Activation in Conventional Therapeutic Exercises and Heavy Resistance Exercises: Implications for Rehabilitation. *Phys Ther*. 2006;86:683-697.

Archer, K.R., Castillo, R.C., MacKenzie, E.J., Bosse, M.J. LEAP study group. Gait symmetry and walking speed analysis following lower-extremity trauma. *Phys Ther*. 2006; 86(12): 1630 – 1640

Baum, M., Nicholas, S., Stock, C., Lee, R. A Proprioceptive neuromuscular facilitation shoulder progression for patients with spinal cord injury resulting in quadriplegia. *Phys Ther Case Reports*. 1998; 1(6): 296 – 300.

Beattie, P.F., Nelson, R.M. Preserving the quality of the patient-therapist relationship: an important consideration for value-centered physical therapy care. *JOPT*. 2008; 38 (2): 34 – 35.

Behram, A., Bowden, M., Nair, P. Neuroplasticity After Spinal Cord Injury and Training: An Emerging Paradigm Shift in Rehabilitation and Walking Recovery. *Phys Ther*. 2006;86:1406-1425.

Bennet J. Informed consent: tips and caveats for PTs. *Liability Awareness*.2007;38-44.

Bleakley CM, McDonough SM, MacAuley DC. Cryotherapy for acute ankle sprains: a randomised controlled study of two different icing protocols. *Br J Sports Med*. 2006;40:700-705. [PNF I has a section on the use of ice in rehabilitation]

Brown LP, Yavorsky P. Locomotor biomechanics and pathomechanics: a review. *JOSPT*. 1987;9:3-10.

Ce E, Margonato V, Casasco M, Veicsteinas A. Effects of stretching on maximal anaerobic power: the roles of active and passive warm-ups. *J Strength Cond Res*. 2008;22(3):794-800.

Cholewicki J, et. al. Delayed trunk muscle reflex responses increase the risk of low back injuries. *Spine*. 2005;30(23):2614-20.

Cholewicki J, Panjabi MM, Khachatryan A. Stabilizing function of trunk flexor-extensor muscles around a neural spine posture. *Spine*. 1997;22:2207-12.

Claësson B. Proprioceptive neuromuscular facilitation (PNF): a literature review. Presented at: IPNFA Conference; September 1996; Göteborg, Sweden.

Cools AM, Witvrouw EE, Declercq GA, et al. Scapular Muscle Recruitment Patterns: Trapezius Muscle Latency With and Without Impingement Symptoms. *Am J Sports Med*. 2003;31:542-549.

Davis DS, Ashby PE, McCale KL, McQuain JA, Wine M. The effectiveness of 3 stretching techniques on hamstring flexibility using consistent stretching parameters. *J Strength and Conditioning Research*. 2005;19(1):27-32.

Dickstein, R., Deutsch, J.E. Motor Imagery in Physical Therapist Practice. *Phys Ther*. 2007; 87 (7): 942 – 953

Drnach, M & Grayhem, P. Striking A Balance: Assessing balance and coordination impairments. *Phys Ther Products*; 2004: 10-15.

Edwards I, Jones M, Hillier S. The interpretation of experience and its relationship to body movement: a clinical reasoning perspective. *Manual Ther*. 2006;11:2-10.

Edin BB. Quantitative analysis of static strain sensitivity in human mechanoreceptors from hairy skin. *J.Neurophysiology*. 1992;67:1105-13.

Edin BB and Johansson N. Skin strain patterns provide kinesthetic information to the human central nervous system. *J.Neurophysiology* 1995;487 ( Pt 1):243-51.

Edin, B.B., Essick, G.K., Trulsson, M., Olsson, K.A., “Receptor encoding of moving tactile stimuli in humans. Temporal Pattern of Discharge of individual low-threshold mechanoreceptors”, *Journal of Physiotherapy*. 1995;15:830-847.

Edin BB and Vallbo AB. Classification of human muscle stretch receptor afferents: a Bayesian approach. *J Neurophysiol.* 1990;63:1314-22.

Edin BB and Vallbo AB. Muscle afferent responses to isometric contractions and relaxations in humans. *J Neurophysiol.* 1990;63:1307-13.

Edin BB and Vallbo AB. Dynamic response of human muscle spindle afferents to stretch. *J Neurophysiol.* 1990;63:1297-306.

Edin BB. The 'initial burst' of human primary muscle spindle afferents has at least two components. *Acta Physiol Scand.* 1991;143:169-75.

Emanuel M, Jarus T, Bart, O. Effect of focus of attention and age on motor acquisition, retention, and transfer: a randomized trial. *Phys Ther.* 2008; 88 (2): 251 – 260.

Eng, JJ, Tang, PF. Gait training strategies to optimize walking ability in people with stroke: a synthesis of the evidence. 2007; 7 (10): 1417 – 1436.

Ennis, M.K. Rewiring the body: neuroplasticity has far-reaching implications for the O&P profession. *Adv Directors in Rehab.* 2007; 43 – 45.

Etnyre B, Thomas DQ. Event Standardization of Sit-to-Stand Movements. *Phys Ther.* 2007;87:1651-1668.

Farrokhi S, Pollard CD, Souza RB, Chen YJ, Rieschl S, Powers CM. Trunk position influences the kinematics, kinetics, and muscle activity of the lead lower extremity during the forward lunge exercise. *JOSPT*, 2008 July; 38

Farthing JP, Borowsky R, Chilibeck PD, Binsted G, Sarty GE. Neuro-physiological adaptations associated with cross-education of strength. *Brain Topography.* 2007;20(2):77-88.(abstract).

Ferber R, Osternig LR, Gravelle DC. Effect of PNF stretch techniques on knee flexor muscle EMG activity in older adults. *J Electromyography and Kine.* 2002;12:391-397.

Folpp H, Deall S, Harvey LA, Gwinn T. Can apparent increases in muscle extensibility with regular stretch be explained by changes in tolerance to stretch. *Aust J Physiother.* 2006;52:45-50.

Friberg R, Thurmond S. Facilitation of the lumbar multifidi and erector spinae using prolonged isometric contraction. Friberg R, Thurmond S. Construct validity of lumbar spine classification system. AAOMPT Conference 2005. (Poster presentation)

Ginn, K., Cohen, M. Exercise Therapy for Shoulder Pain Aimed at Restoring Neuromuscular Control: A Randomized Comparative Clinical Trial. *J of Rehab Med.* 2005;37:115- 122.

Godges JJ, Mattson-Bell M, Thorpe D, Shah D. The immediate effects of soft tissue mobilization with proprioceptive neuromuscular facilitation on glenohumeral external rotation and overhead reach. *J Orthop Sports Phys Ther.* 2003;33:713-718.

Gregory DE, Brown SH, Callaghan JP. Trunk muscle responses to suddenly applied loads: do individuals who develop discomfort during prolonged standing respond differently? *J Electromyogr Kinesiol.* 2008;18(3):495-502.

Grzebellus M., Hering Gernot. “The Effect of contralateral PNF Patterns on Patients After Knee Surgery.”

Hight A.B., Duncan P.W., Nelson S.G. “Electromyographic Activity of Two Contralateral Lower Extremity Muscles During a PNF Pattern.”

Holt, K.G., Saltzman, E., Ho, C.L., Ulrich, B.D. Scaling of dynamics in the earliest stages of walking. *Phys Ther.* 2007; 87 (11): 1458 – 1466

Jette, D.U., Ardleigh, K., Chandler, K., McShea, L. Decision-making ability of physical therapists: physical therapy intervention or medical referral. *Phys Ther.* 2006; 86 (12): 1619 – 1628

Johnson. G., Johnson, V. “The Application of the Principles and Procedures of PNF for the Care of Lumbar Spinal Instabilities.” *J of Manual and Manipulative Ther.* 2002;10:83-105.

Kavcic N, Grenier S, McGill SM. Determining the stabilizing role of individual torso muscles during rehabilitation exercises. *Spine*, 2004 June; 29(11):1254-65

Knott M. In the Groove. *Phys Ther.* 1973;53:365-372. (The Mary McMillan Lecture).

Kofotolis N., Vrabas I.S., Vamvakoudis E., Papanikolaou A., Mandroukas K. Proprioceptive neuromuscular facilitation training induced alterations in muscle fibre type and cross sectional area. *Br J Sports Med.* 2005;39:e11.

Kollen, B., Kwakkel, G., Lindeman, E. Time dependency of walking classification in stroke. *Phys Ther.* 2006; 86 (5): 618 – 625.

Kotoftolis N, Kellis E. Effects of two 4-week proprioceptive neuromuscular facilitation programs on muscle endurance, flexibility, and functional performance in women with chronic low back pain. *Phys Ther.* 2006;86:1001-1012.

- Leerar PJ, Boissonnault W, Domholdt E, Roddey T. Documentation of red flags by physical therapists for patients with low back pain. *J Manual and Manipulative Ther.* 2007;15:42-49.
- Levangie PK, Cook Humphrey E. The shoulder girdle: kinesiology review. *PT Magazine.* December 2000. 48-62.
- Lindemann U, Rupp K, Muche R, Nikolaus T, Becker C. Improving balance by improving motor skills. *Z Gerontol Geriatr,* 2004 Feb; 37
- Ludewig PM, Reynolds JF. The association of scapular kinematics and glenohumeral joint pathologies. *JOSPT.* 2009;39(2):90-104.
- Marchetti, G.F., Whitney, S.L. Construction and validation of the 4-item dynamic gait index. *Phys Ther.* 2006; 86 (12): 1651 – 1667.
- Marchetti GF, Whitney SL, Blatt PJ, Morris LO, Vance JM. Temporal and spatial characteristics of gait during performance of the Dynamic Gait Index in people with and people without balance or vestibular disorders. *Phys Ther.* 2008;88(5):640-651.
- Marek, S.M., Cramer, J.T., Fincher, A.L., Massey, L.L., Dangelmaier, S.M., Purkayastha, S., Fitz, K.A., Culbertson, J.Y. Acute Effects of Static and Proprioceptive Neuromuscular Facilitation Stretching on Muscle Strength and Power Output. *Jour of Athl Train.* 2005; 40(2): 94 - 103.
- Marek SM, Cramer JT, Fincher AL et al. Acute effects of static and proprioceptive neuromuscular facilitation stretching on muscle strength and power output. *J Athletic Training.* 2005;40:94-103.
- McClure, P., Michener, L., Karduna, A. Shoulder Function and 3-Dimensional Scapular Kinematics in People With and Without Shoulder Impingement Syndrome. *Phys Ther.* 2006;86:1075-1090.
- McGill M, Cholewicki J. Biomechanical basis for stability: an explanation to enhance clinical utility. *JOSPT.* 2001;31(2):96-100.
- Moor MA, Kukulka CG. Depression of Hoffman reflexes following voluntary contraction and implications for proprioceptive neuromuscular facilitation therapy. *Phys Ther.* 1991;71:321-333.
- Morris, M. Locomotor Training in People with Parkinson Disease. *Phys Ther.* 2006;86:1425-1435.

Nakamura R, Kosaka K. Effect of proprioceptive neuromuscular facilitation on EEG activation induced by facilitating position in patients with spinocerebellar degeneration. *Tohoku J Exp Med.* 1986;148:159-161.

O'Sullivan P, Dankaerts W, Burnett A, Straker L, Bargon G, Moloney N, Perry M, Tsang S. Lumbopelvic kinematics and trunk muscle activity during sitting on stable and unstable surfaces. *JOSPT*, 2006 Jan; 36

O'Sullivan PB, Twomey L, Allison GT. Dysfunction of the neuro-muscular system in the presence of low back pain: implications for physical therapy. *J Manual and Manipulative Ther.* 1997;5:20-26.

Rundquist, P. Alterations in Scapular Kinematics in Subject with Idiopathic Loss of Shoulder Range of Motion. *JOSPT.* 2007;37:19-25

Schenkman M, Deutsch JE, Gill-Body KM. An integrated framework for decision making in neurologic physical therapist practice. *Phys Ther.* 2006;86(12):1681-1702.

Spernoga SG, Uhl TL, Arnold BL, Gansneder BM. Duration of maintained hamstring flexibility after a one-time, modified hold-relax stretching protocol. *J Athletic Training.* 2001;36:44-48.

Stevenson J, Maitland M, Anemaet W, Beckstead J. Body weight support treadmill training compared with PNF training in persons with chronic spine. Abstract published in *J of Neuro Phys Ther* Dec 2004; presented at APTA combined sections meeting in Feb 2005.

Tsao H, Hodges PW. Persistence of improvements in postural strategies following motor control training in people with recurrent low back pain. *J Electromyogr Kinesiol.* 2007

VanVliet PM, Heneghan NR. Motor control and the management of musculoskeletal dysfunction. *Man Ther*, 2006;11:208-213

Walker, M.L., Austin, A.G., Banke, G.M., Foxx, S.R., Gaetano, L., Gardner, L.A., McElhiney, J., Morris, K., Penn, L. Reference group data for the functional gait assessment. *Phys Ther.* 2007; 87:1468 – 1477.

Wang R. Effect of proprioceptive neuromuscular facilitation on the gait of patients with hemiplegia of long and short duration. *Phys Ther.* 1994;74:1108-1115.

Yekutieli M. The role of vertebral movement in gait: implications for manual therapy. *J Manual and Manip Ther.* 1994;2:22-27.

Yilnen J, Takala EP, Kautiainen H, Nykanen M, Hakkinen A, Pohjolainen T, Karppi SL,

Airaksinen O. Effect of long term neck muscle training on pressure pain threshold: a RCT. *Eur J Pain*, 2005 Dec; 9(6):673-81

Ylinen J, Salo P, Nykanen M, Kautiainen H, Hakkinen A. Decreased isometric neck strength in women with chronic neck pain and the repeatability of neck strength measurements. *Arch Phys Med Rehabil*. 2004;85:1303-1308.